

EDITORIAL OCTOBER 2015

PERACUTE DISEASES WITH MAGIC RESULTS

CASE 1

I had once a chance to see a lady who had gushing menses due to uterine polyp since last 10 days, she was 42 years old, the bleeding was bright red continuous, it was worse any motion like walking, doing simple house hold task, she felt as if she will faint.

She was apparently alright till the time she danced excessively for her son's birthday party, next day she started her menstruation and the bleeding started, she was on a very high dose of Tranexamic acid but she preferred homoeopathy.

Based on her history I selected Erigeron Canadense 1M every few hours, within few hours the bleeding became minimum.

USES IN MY PRACTICE

This remedy is indicated in the management of haemorrhages, when the blood is bright red, also when there is a discharge of mucus.

- It should be remembered in post-partum haemorrhages, also in metrorrhagia, with prolapsus of the uterus.
- The haemorrhage is profuse and a bright red colour.
- The flow is increased by movement, EXERTION ETC.
- It should be studied in epistaxis, and the bleeding from the socket of an extracted tooth, when it may be applied locally and administered internally; also haemoptysis and haematemesis.
- It is an important remedy in cholera infantum, chronic diarrhoea and dysentery, and leucorrhoea when the discharges consist of much mucus mixed with the blood.

CASE 2

A case of cancer of bladder aged 72 years presented in my clinic with severe haematuria, It was T3 N2M1 was already treated with Immunotherapy by intravesicular delivery of Bacillus Calmette–Guérin (BCG) .Here are the symptoms

- severe constipation
- retention of urine very painful
- scanty urine
- urging to urine very violent
- bloody urine in spurts
- feeble stream of urine
- colour dark brown
- offensive urine

Based on above symptoms I selected Arnica 1M few doses which arrested the haemorrhage.

USES IN MY PRACTICE FOR HAEMORRHAGE

- MIND - DELIRIUM - hemorrhage - after
- HEAD - CEREBRAL HEMORRHAGE
- EYE - BLEEDING from eyes - Retinal hemorrhage
- RECTUM - HEMORRHAGE from anus
- URETHRA - HEMORRHAGE
- FEMALE GENITALIA/SEX - METRORRHAGIA - delivery - prevents hemorrhagia
- CHEST - HEMORRHAGE of lungs
- CHEST - HEMORRHAGE of lungs - coagulated
- CHEST - HEMORRHAGE of lungs - coagulated - dark
- CHEST - HEMORRHAGE of lungs - delivery - during
- CHEST - HEMORRHAGE of lungs - exertion agg.; after
- CHEST - HEMORRHAGE of lungs - frothy, foaming
- GENERALS - HEMORRHAGE

- GENERALS - HEMORRHAGE - blood - black
- GENERALS - HEMORRHAGE - blood - bright red
- GENERALS - HEMORRHAGE - blood - clots
- GENERALS - HEMORRHAGE - blood - dark

- GENERALS - HEMORRHAGE - blood - watery

- GENERALS - HEMORRHAGE - Internally

OTHER INDICATIONS

Blunt injuries with extravasation of blood; bruised, sore feeling.

- Mental injuries.
- Overuse of any organ.
- Oversensitive to pain, to being touched.
- Shock from injury.
- * Says he is well when being ill.
- Aversion company.
- Fear of heart-disease.
- * Epistaxis washing face.
- Painful acne.
- Diarrhoea, flatus and odor from mouth smelling like rotten eggs.
- Involuntary urination when running.
- Cough from crying.
- Whooping cough, child cries before coughing.
- Angina pectoris with pain in left elbow.
- Symmetrical eruptions (e.g. left and right hand).
- Black, blue skin.
- Small boils.
- Upper half of body hot, lower cold.
- Heat head with coldness body.

- Bed seems too hard.
- Rheumatism, gout.
- Sprains.
- **Agg.:** cold; wet; touch.
- **Amel.:** lying down with head low.