EDITORIAL FOR JUNE 2017

MY EXPERIENCES WITH SOME MOTHER TINCTURES LOCALLY AS WELL AS INTERNALLY

I will begin with external application of mother tincture even though Hahnemann has given strictest warning regarding the use of local application in his chronic disease. I somehow feel this is truer for skin diseases and venereal diseases and hence I freely use mother tinctures as a local application in cases of injuries, wounds, gangrene, post-surgical complications. Here are some of the quotes from Hahnemann from chronic disease

"Allopathic doctor destroys in his blindness, through local applications"

"But if the chancre is destroyed through local applications * before the internal disease is healed, -and this is still a daily practice with physicians of the old school."

"Just as emphatically Fabre declares: ++ "Syphilis always follows on the destruction of the chancre by local applications"

"Psora through anti-psoric remedies is effected most easily only while the original eruption of itch is still present. From this it again appears how unconscionable it is of the allopathic physicians, to destroy the primitive itch eruption through local applications"

The following medicines given below has been used by me very successfully without the trace of suppression as claimed by Hahnemann. There is nothing wrong with clinical experiments if one can draw honest conclusions

Anthracinum

- If the swelling changes colour and red streaks from the wound map out the course of lymphatics
- Suspicious insect stings.
- Tendency is to become gangrenous;
- Wounds from accidents where there is indication of threatened gangrene.

Calendula officinalis

- Clean, surgical cuts or lacerated wounds
- External wounds with or without loss of substance; torn and jagged looking wounds;
- Lacerated wounds
- Post-surgical operations; to promote healthy granulation and prevent excessive suppuration and disfiguring scars.
- Rupture of muscles or tendons; lacerations during labour;
- Traumatic and idiopathic neuroma
- Wounds penetrating articulations with loss of synovial fluids.

Formalinum

• Eczema in neighbourhood of wound.

Hypericum perforatum

• Punctured, incised or lacerated wounds;

Ledum palustre

- Punctured wounds by sharp-pointed instruments, as awls, nails
- Rat bites, stings of insects, especially mosquitoes.

Liatris spicata

• Locally, applied to ulcers and unhealthy wounds.

Sempervivum tectorum

• Locally for bites of insect, stings of bees, and poisoned wounds.

Dosage

All local application should be diluted 50% with distil water. The standard doses to apply 4 times a day with a clean sterilized gauze piece

Mother tinctures which I used internally I very rarely use mother tinctures over the period of years it's use in my practise has been diminished considerably, the main reason is they cannot cure as the dose is not dynamic. However, I will discuss few which has been used successfully

Fraxinus americans

• Fibroid uterus

Galium aparine

• Homeopathic diuretic

Hygrophilia spinose

• Homeopathic diuretic

Liatris spicata

• Homeopathic diuretic due to liver and renal disease

Nitri spiritus dulcis

• Homeopathic diuretic

Oxydendron

• Homeopathic diuretic

Ruta

• Nodes, nodules in tendons and muscles

Scrophularia nodosa

• Fibroadenoma of the breast

Solidago v

• Epithelial cells in the urine

Thalapsi bursa pastoris

- Dysuria in old person
- Endometriosis

Dosage

2 drops in 1 teaspoonful of water 3 times a day.

Warning

Do not use mother tinctures as routine. This is not at all classical homeopathy. This will never cure the disease it will only palliate